



FLORIST

You and your medicine

This booklet does not replace the Patient Information Leaflet (PIL) included in your medicinal pack. Please refer to the PIL for full information.

Vosevi is indicated for the treatment of chronic hepatitis C virus (HCV) infection in patients aged 12 years and older and weighing at least 30 kg.<sup>1,2</sup>

The dose referred to in this patient booklet is for adults only. This material should only be given to adult patients.

**Reporting of side effects:** If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report any side effects you may get directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. For Ireland, Reporting forms and information can be found at www.hpra.ie and can be reported to HPRA on +35316764971. By reporting side effects, you can help provide more information on the safety of this medicine.

UK-VSV-0087 | Date of preparation: March 2024



This material has been created, distributed and funded by Gilead Sciences Ltd.

### **Contents**

I've been diagnosed with hepatitis C (hep How could this medicine help me? Is this medicine suitable for me? What if I miss a tablet? What if I drink alcohol or take illicit drugs v Can I take other medicines with Vosevi? What about pregnancy or breastfeeding? Can I be infected with hep C again? You're not on your own Talk openly with your healthcare team References



C) 04	
05	5
Oe	5
10	)
while on my Vosevi medicine? <b>1</b> 1	
12	
14	
15	;
16	5
17	,
18	}

# I've been diagnosed with hepatitis C

Hepatitis C (hep C) is a virus that can infect the liver.<sup>3</sup> The hepatitis C virus lives in infected blood (and other bodily fluids) and is most commonly spread when someone comes into contact with blood from an infected person.<sup>3,4</sup> Your diagnosis of hep C may have you feeling worried, but diagnosis is the first step in the journey to improving your health.

# How could this medicine help me?

Your healthcare provider has given you Vosevi to treat hepatitis C. As long as the medicine is taken in the recommended way, more than 90% of people **may be** cured.<sup>1-3,5,6</sup>

### What is the medicine I've been given?

Vosevi is a beige, capsule-shaped, film-coated tablet containing substances called sofosbuvir, velpatasvir and voxilaprevir. On one side it is stamped with the letters "GSI" and on the other side is the number "3".<sup>12</sup>



Medicines for hep C can sometimes cause side effects – headache, diarrhoea or feeling sick. Your healthcare team should be able to suggest things to help ease any discomfort.<sup>1,2,6</sup>

Not to scale.



#### of patients with hep C may be cured.<sup>1,2,5</sup>



Take one tablet, once a day for 8–12 weeks to help treat the virus, depending on how your doctor has advised you to take it.<sup>1.2</sup>

# Is this medicine suitable for me?

You and your healthcare team will have discussed the most suitable treatment options for you. This might have included a discussion about your lifestyle as well as medicines that may suit you. If you have been given Vosevi, your healthcare team think it's an appropriate medicine for you.

Remember to tell your healthcare team about anything else you are taking - whether that's medicines, herbal remedies or illicit drugs.<sup>2,7</sup>

If you have any questions about your medicines, please ask your healthcare team.

If you're in a sexual relationship you'll need some effective contraception for you and your partner while you're being treated for your hepatitis C infection, so ask for something suitable. This will minimise the risk of passing it on to your partner or an unborn child.

You should also tell your healthcare team if any of the following apply to you before you take your Vosevi medicine:<sup>2,7</sup>

- You have problems with your liver or you have had a liver transplant
- You are living with hepatitis B, or have had a previous hepatitis B infection
- You have kidney problems or if you are on kidney dialysis
- You're taking other medicines, herbal remedies or illicit drugs
- You're pregnant and/or breastfeeding
- You're living with HIV
- You're taking, or have recently taken, any medications for heart conditions
- You have diabetes

### How do I take my Vosevi medicine?

•

Take one tablet, once a day, with food. The tablet should not be crushed, split, or dissolved, so make sure you swallow it whole.<sup>1,2</sup> For the best chance of cure, it is important to take your medicine as instructed by your doctor.



If, however, your doctor has prescribed you another medicine called a proton pump inhibitor (such as omeprazole, lansoprazole, rabeprazole, pantoprazole or esomeprazole), and you need high doses, your doctor may give you a different medicine or adjust the dose of the medicine you're taking.<sup>12,7</sup>

If you are taking an antacid, take it at least 4 hours before or at least 4 hours after Vosevi.2,7

If your doctor has prescribed an H2-antagonist (such as famotidine, cimetidine, nizatidine or ranitidine), and you need high doses, your doctor may give you a different medicine or adjust the dose of the medicine you're taking.<sup>2,7</sup>

Your healthcare team may ask you to come back in for a check-up after 12 or 24 weeks to check that the virus has been cleared from your body. They'll carry out a blood test to do this. In a small number of people the virus is not cleared after treatment. If you're one of these people, your healthcare team may suggest you try another medicine.<sup>2,6,7</sup>

Remember to read the Patient Information Leaflet for more details on how to take your medicine.



# How should I store my Vosevi medicine?

There are no special storage requirements for Vosevi, so no need to keep it in the fridge. Your medicine should be kept out of sight and reach of children and animals, and should not be used after the expiration date shown on the packaging.<sup>2,7</sup>

If the seal is broken or missing when you first open your Vosevi medicine, tell your healthcare team immediately. Always dispose of any unused medicine through your pharmacist. Do not share your medicine with anyone else.<sup>2,7</sup>

## What if I miss a tablet?

Try to take your medicine at the same time every day – it will help you to remember. For example, when you get up in the morning or when you clean your teeth.

It's very important that you complete the full course of treatment to give the medicine the best chance of working to treat the hep C virus infection. If you do forget and miss taking your medicine at any point during the 8–12 weeks, it's not ideal but try not to worry – the information below can help you continue your treatment for the best chance to clear the virus.

First of all, work out how long it is since you last took a tablet.

### 18 hours

If you notice **within 18 hours** of missing your dose, it's ok to take the tablet you missed straight away.<sup>27</sup> Take your next tablet at your usual time and carry on as before.<sup>1.27</sup>

If it's been **more than 18 hours** since you last took your Vosevi tablet, just miss that dose out. *Simply take your next dose as normal.*<sup>1,2,7</sup>



It's important that you don't take a double dose (that's two doses close together in time) so try to keep track.<sup>12,7</sup>

If you've been distracted recently and have missed more than one dose, please inform your healthcare team.

What if I drink alcohol or take illicit drugs while on my Vosevi medicine?





If you think that you will take illicit drugs or drink alcohol while taking your medicine, it's important that you tell your healthcare team. They can support you, and are there to give you help and advice without judgement.



#### Can I take other medicines with Vosevi?

If you're taking other medicines including any herbal remedies (such as St John's Wort), you must tell your healthcare team.<sup>2.7</sup> This is because some things can interact with your medicine and may make it less effective or work in a different way. It's important to mention everything – even if you only take it occasionally. This also includes discussing any contraception you might be taking particularly those including ethinylestradiol.<sup>2,7</sup> If you're in a sexual relationship, you should not risk passing the infection on to your partner or an unborn child before you're cured, so you'll need effective contraception while you're being treated for hepatitis C.<sup>1,2,7</sup>





#### What about pregnancy or breastfeeding?

Vosevi is not recommended for use during pregnancy. This is because we don't know much about how Vosevi might affect pregnant women.<sup>1,2</sup>

It is important not to breastfeed during treatment, as it is not known if some of the active substances of Vosevi may pass into human breast milk.<sup>2,7</sup>

If you or your partner become pregnant during treatment with Vosevi, please contact your doctor or healthcare team immediately.<sup>2,7</sup>



#### I'm worried about side effects


Like any other medicine, there are some common side effects with Vosevi such as headache, diarrhoea, feeling sick, stomach pain, decreased appetite, being sick, muscle pain, abnormal liver function tests and rash. Stop taking Vosevi and seek medical help immediately if you experience swelling of the face, lips, tongue or throat or if you develop a wide-spread severe rash with peeling skin, which may be accompanied by fever, flu like symptoms, blisters in the mouth, eyes, and/or genitals (Stevens Johnson syndrome).<sup>2,7</sup> If you notice any side effects or you have any other questions about taking your medicine, just ask your healthcare provider. They'll be very happy to answer your guestions. Remember to read the Patient Information Leaflet for more details on how to take your medicine and how to report any adverse events.

You can still spread hep C during your treatment, so it's important to take your Vosevi medicine for the whole length of treatment. Successfully treating hep C means that you cannot pass it on to other people anymore.<sup>3</sup> But even after you have finished treatment, you can catch hep C again.<sup>3</sup> To reduce the risk of catching hep C again or passing it on, avoid:4

- Sharing needles, syringes and other items used to take drugs
- Having unprotected sex this includes men who have sex with men
- Sharing your razor or toothbrush

Can I transmit my infection while on treatment and be infected with hep C after being cured?

 $\mathbf{UO}_{\mathbf{O}}$ 





If you have other thoughts or questions that aren't answered by this booklet, the best place to start is with your healthcare team - they are always ready to help you. Try to be as open as you can with them so they can help you in the best way possible. If you'd like to ask some questions, but can't think of any, here are some common ones that could help you get started. When starting treatment How do I take the treatment? What do I need to know during treatment? Is there anything I need to tell my healthcare team How often do I need to see my healthcare team, before taking my medicine? and will they arrange the appointments for me? How long before I'm cured? How can I avoid getting hepatitis C again? After treatment What support services are available to me and how do I get in touch with them?

### References

### Notes

1.	Vosevi 400 mg/100 mg/100 mg film coated tablets. Summary of Product Characteristics, UK.
	[Online] Available at: https://www.medicines.org.uk/emc/product/772/smpc.

- Vosevi 400 mg/100 mg/100 mg film coated tablets. Summary of Product Characteristics and Package Leaflet, Ireland. [Online] Available at: https://www.ema.europa.eu/en/documents/product-information/vosevi-epar-product-information\_en.pdf.
- 3. NHS. Conditions. Hepatitis C. Overview. [Online] Available at: https://www.nhs.uk/conditions/hepatitis-c/.
- NHS. Conditions. Hepatitis C. Causes. [Online] Available at: https://www.nhs.uk/conditions/hepatitis-c/causes/.
- 5. World Health Organization. Hepatitis C Factsheet.. [Online] Available at: https://www.who.int/news-room/fact-sheets/detail/hepatitis-c.
- 6. NHS. Conditions. Hepatitis C. Treatment. [Online] Available at: https://www.nhs.uk/conditions/hepatitis-c/treatment/.
- 7. Vosevi 400 mg/100 mg/100 mg film coated tablets. Patient Information Leaflet, UK. [Online] Available at: https://www.medicines.org.uk/emc/product/772/pil.



Notes		Im	



